

Coisotropic D-branes on $\text{AdS}_4 \times \mathbb{CP}^3$ and massive deformations

Paul Koerber

*Max-Planck-Institut für Physik
Föhringer Ring 6, D-80805 München, Germany*

E-mail: koerber at mppmu.mpg.de

ABSTRACT: We scan for massive type IIA $\text{SU}(3)$ -structure compactifications of the type $\text{AdS}_4 \times \mathbb{CP}^3$ with internal symmetry group $\text{SO}(4)$. This group acts on $\mathbb{CP}^3$ with cohomogeneity one, so that one would expect new non-homogeneous solutions. We find however that all such solutions enhance their symmetry group to $\text{Sp}(2)$ and form, in fact, the homogeneous family first described in [1]. This is in accordance with [2], which argues from the CFT-side that although new vacua with $\text{SO}(4)$ symmetry group and $N = 2$ supersymmetry should exist, they fall outside our ansatz of strict $\text{SU}(3)$ -structure, and instead have genuine $\text{SU}(3) \times \text{SU}(3)$ -structure. We do find that the $\text{SO}(4)$ -invariant description, which singles out one preferential direction in the internal space, is well-adapted for describing the embedding of AdS_4 -filling supersymmetric D8-branes on both the original ABJM configuration as its massive $\text{Sp}(2)$ -symmetric deformations. Supersymmetry requires these D-branes to be of the coisotropic type, which means in particular that their world-volume gauge field must be non-trivial.

Contents

1. Introduction	1
2. Setup: $\text{SO}(4)$-invariant ansatz	4
2.1 Symmetry group and codimension-one foliation	4
2.2 The coset $T^{1,-1}$	7
2.3 Starting point: the massless Kähler-Einstein geometry	9
3. Supersymmetry conditions and Bianchi identities	11
3.1 Review of the conditions for supersymmetric AdS_4 solutions	11
3.2 The massless Kähler-Einstein geometry revisited	13
3.3 Solving the conditions	14
3.4 Equivalence to the family of $N = 1$ massive $\text{Sp}(2)$ -invariant solutions	17
4. Coisotropic D-brane embeddings	19
5. Concluding remarks	21

1. Introduction

Supersymmetric compactifications with fluxes have become an important ingredient in the search for realistic vacua of string theory, since some or even all of the moduli can be stabilized (for reviews see e.g. [3, 4, 5, 6]). Apart from phenomenological reasons, the philosophy behind imposing supersymmetry is that solving for the supersymmetry conditions is much easier than solving all the equations of motion, while it has been shown that for (eleven-dimensional and type II) supergravity these supersymmetry conditions completed with the Bianchi identities indeed imply the equations of motion [7, 8, 9, 10]. To obtain solutions with a positive cosmological constant, however, one has to break supersymmetry. A further complication for constructing compactifications with a non-negative four-dimensional cosmological constant is that in the presence of fluxes the no-go theorem of [11] requires the introduction of negative-tension sources. Although in string theory they can be provided for by orientifolds, these localized sources complicate the explicit form of the solution. The strategy is then to construct solutions with a negative cosmological constant, with AdS_4 as the external space, and take care of both uplifting to dS_4 and supersymmetry breaking in a later stage.

Because of the gauge/gravity correspondence type IIA solutions with an AdS_4 factor are also interesting as geometries potentially dual to a three-dimensional conformal theory. Following [12], where the three-dimensional CFT is a Chern-Simons-matter theory, there

has been lot of progress in this direction recently. In this way these solutions can inspire the construction of new three-dimensional CFTs or, conversely, the CFTs can lead the way to the construction of new geometries.

The supersymmetry and Bianchi conditions for type IIA AdS_4 flux solutions with strict $\text{SU}(3)$ -structure were first worked out in [8].¹ They lead to very specific requirements on the internal geometry. In particular, only the $\text{SU}(3)$ -structure torsion classes $\mathcal{W}_1$ and $\mathcal{W}_2$ can be non-zero and they can be chosen to be purely imaginary.

Prominent examples of solutions to these equations are homogeneous constructions on coset manifolds [13, 14, 15, 16, 1, 17]. In particular, in [1] a family with one shape parameter was constructed on $\mathbb{CP}^3 = \frac{\text{Sp}(2)}{\text{S}(\text{U}(2) \times \text{U}(1))}$. Changing the shape parameter, these $\text{Sp}(2)$ -invariant solutions interpolate between the solution of [18] – with vanishing Romans mass and standard Fubini-Study metric – and a second, squashed, massless solution. While the two endpoints of the family can be lifted to M-theory and correspond to the standard S^7 and the squashed S^7 -solution respectively, the rest of the family has non-zero Romans mass and includes in particular a nearly-Kähler solution (i.e. $\mathcal{W}_2 = 0$). It was argued that a similar story applies to $\frac{\text{SU}(3)}{\text{U}(1) \times \text{U}(1)}$, although it was later found [17] that these $\text{SU}(3)$ -invariant solutions actually have two shape parameters. The effective four-dimensional theory corresponding to an expansion in terms of left-invariant forms on these cosets was studied in [19, 20]. It was found in [21] that this expansion actually corresponds to a consistent reduction.

The generic solution in the family has $N = 1$ supersymmetry. However, for the massless solution on $\mathbb{CP}^3$ with Fubini-Study metric, the supersymmetry enhances to $N = 6$. At the same time the bosonic symmetry group enhances from $\text{Sp}(2)$ to $\text{SU}(4)$ and corresponds to the description of $\mathbb{CP}^3$ as the coset manifold $\frac{\text{SU}(4)}{\text{S}(\text{U}(3) \times \text{U}(1))}$. It got recently a lot of attention as the dual geometry for the original ABJM gauge/gravity correspondence [12] in the limit where the type IIA description is valid. The geometry then also has a Kähler form and becomes Einstein, so we will dub it the massless Kähler-Einstein solution in the following. Also for the squashed massless case, a CFT dual has in the meantime been proposed [22].

A natural question that arises then, namely whether also CFT duals can be found for the interpolating solutions with non-zero Romans mass, was addressed in [2] (see also [23]). Surprisingly, not only these could be found, but in fact several massive deformations of the ABJM Chern-Simons-matter theory were proposed with different numbers of supersymmetries and different global symmetry groups: $N = 0$ with $\text{SO}(6)$ -symmetry, $N = 1$ with $\text{Sp}(2)$ symmetry, $N = 2$ with $\text{SO}(4) \times \text{SO}(2)_R$ and finally $N = 3$ with $\text{SO}(3) \times \text{SO}(3)_R$. It was argued that the $N = 0$ theories correspond to AdS_4 solutions with the standard Fubini-Study metric on $\mathbb{CP}^3$, but with different fluxes (and in particular of course non-zero Romans mass) from the ones in the $N = 6$ configuration. Similar non-supersymmetric AdS_4 vacua were constructed in [24, 25], and it would be interesting to investigate whether they also allow for CFT duals. The second family with $N = 1$ supersymmetry and $\text{Sp}(2)$

¹I use here the terminology *strict* $\text{SU}(3)$ -structure for a supersymmetry ansatz where the same globally-defined internal spinor enters in both the left- and right-moving supersymmetry generator of type II supergravity, in order to distinguish it from the more general $\text{SU}(3) \times \text{SU}(3)$ -structure ansatz, where two different internal spinors enter.

bosonic symmetry group then corresponds to the dual of the above interpolating solutions, while we come back to the last two families in a minute.

In this paper I will scan for supersymmetric solutions on $\mathbb{CP}^3$ with strict $SU(3)$ -structure and a $SO(4)$ symmetry group, embedded in $SU(4)$ such that it acts with cohomogeneity one on $\mathbb{CP}^3$.² With only one transversal coordinate this would provide the easiest setup for constructing non-homogeneous solutions. Unfortunately, we find that all such solutions enhance their symmetry group to $Sp(2)$ and are in fact the known family of $N = 1$ solutions mentioned before, in disguise.

This is in agreement with the results of [2], where although the third family of three-dimensional Chern-Simons-matter theories presented there has $SO(4)$ global symmetry, it also has $N = 2$ supersymmetry and $SO(2)$ R-symmetry. It was then argued that the geometric dual to this CFT could not be described with a strict $SU(3)$ -structure ansatz, but would instead have two genuine $SU(3) \times SU(3)$ -structures. The reason is that the supersymmetry conditions of [8] directly relate the $(3,0)$ -form Ω with the NSNS-flux H . So if both supersymmetries would be of the strict $SU(3)$ -structure type, they would correspond to the same Ω and thus the same almost complex structure. Since from the metric and the almost complex structure one can then also find the two-form J , the $SU(3)$ -structures and thus supersymmetries would in the end be entirely the same. Furthermore, since the R-symmetry requires both supersymmetries to be of the same type, they must be both $SU(3) \times SU(3)$. The only way around the argument is for $m = 0$ where also $H = 0$ and there is no relation anymore between Ω and H . This loophole makes it possible for the standard ABJM $\mathbb{CP}^3$ to be $N = 6$ and still have strict $SU(3)$ -structure. In the meantime the geometries with $SU(3) \times SU(3)$ -structure corresponding to these $N = 2$ (and also $N = 3$) CFTs were constructed in [29] to first order in the mass parameter. A similar geometry on $M^{1,1,1}$ was obtained in [30].

The $SO(4)$ -invariant setup is however useful to describe supersymmetric AdS_4 -filling D8-branes. Indeed, these D-branes will have one transversal direction and wrap the five-dimensional internal cycle on which the $SO(4)$ acts homogeneously. According to [31, 32] the supersymmetry conditions for D-branes force them to be generalized calibrated. Just as for the supergravity background, they are also sufficient to ensure that the D-brane solves its equations of motion. In the case of AdS_4 compactifications there are some additional subtleties that were studied in detail in [33]. One interesting aspect is, for instance, that the so-called D-flatness condition will follow automatically from the F-flatness conditions. For space-time-filling D6-branes, for example, that means that a Lagrangian cycle will automatically be special (with respect to the $SU(3)$ -structure (J, Ω) associated to the bulk supersymmetry). For the particular case of space-time-filling D8-

² There is another inequivalent embedding of $SO(4)$ in $SU(4)$, which acts with cohomogeneity three. This is the symmetry group of the Lagrangian D-branes of [26, 27, 28]. The difference can also be seen on how it acts on the $\mathbf{6}$ (the antisymmetric two-tensor) of $SU(4)$, the representation under which the supersymmetry generators transform. Under the first $SO(4) = SU(2) \times SU(2)$ – the one considered in this paper – it decomposes as $\mathbf{6} \rightarrow (\mathbf{1}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{1}) \oplus (\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2})$, and there are two supersymmetry generators singled out corresponding to the two first terms in the decomposition. For the second $SO(4)$ we find $\mathbf{6} \rightarrow (\mathbf{3}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{3})$. The $N = 3$ supersymmetry preserved by a Lagrangian D6-brane correspond then to the first term.

branes, the supersymmetry conditions imply that they must be coisotropic [34], which means in particular they must have a non-trivial world-volume gauge field. Examples of coisotropic D-branes are still quite rare: there are constructions on the torus (see e.g. [35] for an application) and in the context of the world-sheet approach (see e.g. [36]). In the latter case it is however not yet clear whether the D-flatness condition can be satisfied. Coisotropic D-branes (on the Hitchin moduli space) also played role in the construction of [37]. In this paper, I construct coisotropic embeddings on the whole family of $N = 1$ $\text{Sp}(2)$ -invariant solutions. In the Kähler-Einstein limit these coisotropic D-branes will preserve $N = 2$, while away from the limit they preserve the $N = 1$ of the bulk.

In section 2 I study the constraints on the solutions from having the $\text{SO}(4)$ symmetry group with the embedding in $\text{SU}(4)$ discussed above. I find that the solution should have cohomogeneity one, and that the five-dimensional orbits of $\text{SO}(4)$ form the coset $T^{1,-1} = \frac{\text{SO}(4)}{\text{U}(1)}$, better known as the base of the conifold [38]. I rewrite the starting point, namely the $N = 6$ massless solution with Fubini-Study metric, in terms of coordinates reflecting the $\text{SO}(4)$ -symmetry. In section 3.1 I review the conditions for strict $\text{SU}(3)$ -structure of [8], which I solve in section 3.3, starting from the ansatz suggested by the symmetry analysis of section 2. In section 3.4, I show that the single resulting family of solutions is actually equivalent to the known family of $\text{Sp}(2)$ -invariant solutions. In section 4 I construct coisotropic D-brane embeddings on this family.

2. Setup: $\text{SO}(4)$ -invariant ansatz

In this section I analyse the constraints that the $\text{SO}(4)$ symmetry group imposes on the metric and the form fields of the solution. It turns out that the $\text{SO}(4)$ symmetry is not large enough to make the solutions homogeneous. Instead, they have cohomogeneity one, which is the next easiest case. The ansatz for the metric and the form fields will then depend on functions in only one variable.

2.1 Symmetry group and codimension-one foliation

The ABJM theory [12] consists of a $\text{U}(N)_k \times \text{U}(N)_{-k}$ Chern-Simons theory with chiral multiplets

$$(A_1, A_2, B_1^*, B_2^*) \quad (2.1)$$

transforming as a **4** under the global symmetry group $\text{SU}(4)$. According to [2] turning on the Romans mass, in the $N = 2$ branch this global symmetry is broken to $\text{SO}(4) \times \text{SO}(2)_R$. The $\text{SO}(4) = \text{SU}(2)_A \times \text{SU}(2)_B$ part acts such that

$$(A_1, A_2), \quad (B_1, B_2) \quad (2.2)$$

transform as **2_A** and **2_B** respectively.

The dual of the massless ABJM theory in the regime where type IIA is valid, is the $\text{AdS}_4 \times \mathbb{CP}^3$ solution of [18]. The internal space $\mathbb{CP}^3$ is then equipped with the Fubini-Study metric and is described by the coset

$$\frac{\text{SU}(4)}{\text{S}(\text{U}(3) \times \text{U}(1))}. \quad (2.3)$$

The homogenous coordinates of $\mathbb{CP}^3$,

$$(Z^1, Z^2, Z^3, Z^4) \cong \lambda(Z^1, Z^2, Z^3, Z^4), \quad \text{with } \lambda \in \mathbb{C}_*, \quad (2.4)$$

then transform as a **4** just like the (A_i, B_i^*) in eq. (2.1). We therefore deduce that for the deformed geometry the isometry $\text{SO}(4)$ should act such that

$$(Z^1, Z^2), \quad (Z^3, Z^4) \quad (2.5)$$

transform as **2_A** and **2_B** respectively. This particular $\text{SO}(4)$ -embedding thus breaks **4** $\rightarrow$ **(2, 1)** $\oplus$ **(1, 2)** and as we will discuss in section 3.4 also **6** $\rightarrow$ **(1, 1)** $\oplus$ **(1, 1)** $\oplus$ **(2, 1)** $\oplus$ **(1, 2)**. As discussed in footnote 2 and as we will show presently, it is this embedding of $\text{SO}(4)$ that acts with cohomogeneity one on the internal space.

So let us study the orbits of $\text{SO}(4)$. First we turn to special points with coordinates of the form

$$(Z^1, Z^2, 0, 0). \quad (2.6)$$

One can easily check that they are rotated into each other by $SU(2)_A$ and that the part of the original isotropy group $S(U(3) \times U(1))$ that belongs to $\text{SO}(4)$ is $U(1) \times SU(2)_B$. The $SU(2)_B$ factor cancels out in numerator and denominator so that the orbit is

$$\mathbb{CP}_A^1 = \{(Z^1, Z^2) : (Z^1, Z^2) \cong \lambda(Z^1, Z^2)\} = \frac{SU(2)_A}{U(1)}. \quad (2.7)$$

Likewise, the orbit of the points of the special form

$$(0, 0, Z^3, Z^4) \quad (2.8)$$

is also a $\mathbb{CP}^1$, which we call $\mathbb{CP}_B^1$.

Let us now turn to the orbit through a point with coordinates of the form

$$(0, Z^2, Z^3, 0). \quad (2.9)$$

Since acting with $\text{SO}(4)$ one can bring a generic point in this form, this is the generic case. We find that this time the part of the isotropy group $S(U(3) \times U(1))$ that belongs to $\text{SO}(4)$ is a $U(1)$ acting like

$$(Z^1, Z^2, Z^3, Z^4) \rightarrow (e^{i\phi} Z^1, e^{-i\phi} Z^2, e^{-i\phi} Z^3, e^{i\phi} Z^4). \quad (2.10)$$

We conclude that the orbit of $\text{SO}(4)$ through this point is five-dimensional and given by the coset manifold

$$\frac{\text{SO}(4)}{U(1)} = T^{1,-1}, \quad (2.11)$$

which is better known as the base of the conifold (see e.g. [38]). It has the topology of $S^2 \times S^3$. $\mathbb{CP}^3$ can thus locally be considered as a foliation with the generic leaves taking the form of the coset $T^{1,-1}$. This is not globally a foliation though, since this description breaks down at special points of the forms (2.6) and (2.8), where the leaves collapse to two-dimensional $\mathbb{CP}^1$ s. When constructing metrics on $\mathbb{CP}^3$ in this description it is therefore




Figure 1: Representation of the geometry of the solutions on $\mathbb{CP}^3$. As explained in the text, the coset $T^{1,-1}$ can be seen as a $U(1)$ -bundle over $\mathbb{CP}_A^1 \times \mathbb{CP}_B^1$. How the sizes of these $\mathbb{CP}^1$ s vary with t depends on the solution and is described by the metric (2.28) in the Kähler-Einstein case or by the metric (3.32) for the general solution. In all cases the $U(1)$ and $\mathbb{CP}_A^1, \mathbb{CP}_B^1$ shrink to zero for $t \rightarrow 0, +\infty$ respectively.

important to carefully check the regularity on these degenerate orbits. That $\mathbb{CP}^3$ could be locally seen as a foliation in this way was noted before in [39].

Another way to see this foliation, which allows for a better handle on the location of the two special loci, is to start from a generic point (Z^1, Z^2, Z^3, Z^4) and note that the action of $SO(4) = SU(2)_A \times SU(2)_B$ allows to go to any other point with the same absolute values $|Z^1|^2 + |Z^2|^2$ and $|Z^3|^2 + |Z^4|^2$. Since these are homogenous coordinates, there is only meaning to the relative factor $t = (|Z^1|^2 + |Z^2|^2) / (|Z^3|^2 + |Z^4|^2)$. This implies that the orbit of the point (Z^1, Z^2, Z^3, Z^4) contains all the points with the same relative factor t . The two special orbits then correspond to sending the factor t to zero and infinity respectively. The first two coordinates can be seen to describe a $\mathbb{CP}^1 = S^2$, for which we can choose inhomogeneous coordinates fixing the scaling freedom. For the last two coordinates the absolute value $|Z^3|^2 + |Z^4|^2$ is then fixed so that we find an S^3 . In the special loci, the radius of this three-sphere or the one with the roles of (Z^1, Z^2) and (Z^3, Z^4) interchanged goes to zero. See figure 1 for a an illustration of the geometry.

The space $T^{1,-1}$ can also be considered as a $U(1)$ -bundle over $\mathbb{CP}_A^1 \times \mathbb{CP}_B^1$. Indeed the two $\mathbb{CP}^1$ s are described by the homogeneous coordinates (Z^1, Z^2) and (Z^3, Z^4) respectively, and the $U(1)$ describes the relative phase between the two factors.

It will turn out that in the massless Kähler-Einstein limit the $SO(2)_R$ symmetry that

rotates both $\text{SO}(4)$ -invariant supersymmetries into each other, acts as follows

$$(Z^1, Z^2) \rightarrow e^{i\phi/2}(Z^1, Z^2), \quad (Z^3, Z^4) \rightarrow e^{-i\phi/2}(Z^3, Z^4). \quad (2.12)$$

Let us now study the coset $T^{1,-1}$, which makes up the geometry of the five-dimensional orbits, in a bit more detail.

2.2 The coset $T^{1,-1}$

Imposing that our solutions respect the $\text{SO}(4)$ symmetry group, implies that the metric, the $\text{SU}(3)$ -structure and the form fields should all be invariant under the left action of this group. In this section we construct the most general form of the $\text{SO}(4)$ left-invariant metric and forms on the $T^{p,q}$ cosets. For a review on coset manifolds and their description in terms of left-invariant forms see [40, 41]. For an overview of the $T^{p,q}$ coset in particular – which also explains its description as a $\text{U}(1)$ bundle over $\mathbb{CP}^1 \times \mathbb{CP}^1$ – see [38]

As explained in these reviews (and also in [17, 19]) on a coset manifold G/H we must split off the generators H_a of the algebra corresponding to the group H that is projected out from the rest of the generators E_i of G . This induces a split of the structure constants:

$$\begin{aligned} [H_a, H_b] &= f_{ab}^c H_c, \\ [H_a, K_i] &= f_{ai}^j K_j + f_{ai}^b H_b, \\ [K_i, K_j] &= f_{ij}^k K_k + f_{ij}^a H_a, \end{aligned} \quad (2.13)$$

where for a reductive coset, which is the case for us, the split basis can be chosen such that $f_{ai}^b = 0$. The decomposition of the Lie-algebra valued one-form

$$L^{-1}dL = e^i K_i + \omega^a H_a, \quad (2.14)$$

for $L \in G$, defines then a coframe $e^i(y)$ on G/H . The left-invariant l -forms ϕ are then the forms which can be expanded in terms of the coframe e^i with coefficients constant over the coset, and moreover satisfy

$$f_{a[i_1}^j \phi_{i_2 \dots i_l]j} = 0. \quad (2.15)$$

In our case however, we must allow the coefficients to depend on the one coordinate transversal to the coset. Likewise a left-invariant metric must have coefficients constant over the coset in the coframe basis e^i and satisfy the condition

$$f_{a(i}^k g_{j)k} = 0. \quad (2.16)$$

The exterior derivatives of the one-forms of the coframe can be expressed in terms of the structure constants as follows:

$$de^i = -\frac{1}{2}f_{jk}^i e^j \wedge e^k - f_{aj}^i \omega^a \wedge e^j. \quad (2.17)$$

Acting with the exterior derivative on left-invariant forms, the condition (2.15) will ensure that all terms containing ω^a drop out.

Suppose the $SU(2)_{A/B}$ factors in $SO(4)$ are generated by $\lambda_{A,B} = -\frac{i}{2}\sigma_{A,B}$, where $\sigma_{A,B}$ are the Pauli matrices, then we propose the following split for the generators of the algebra of $SO(4)$

$$(E_1, \dots, E_5) = (\lambda_{A1}, \lambda_{A2}, \lambda_{B1}, \lambda_{B2}, \frac{1}{2p}\lambda_{3A} + \frac{1}{2q}\lambda_{3B}),$$

$$H_1 = \frac{1}{2p}\lambda_{3A} - \frac{1}{2q}\lambda_{3B}.$$
(2.18)

Here H is the generator of the $U(1)$ that is projected out.

This leads to the structure constants:

$$f^5_{12} = f^6_{12} = p, \quad f^5_{34} = -f^6_{34} = q, \quad f^1_{25} = f^2_{51} = \frac{1}{2p}, \quad f^3_{45} = f^4_{53} = \frac{1}{2q},$$

$$f^1_{26} = f^2_{61} = \frac{1}{2p}, \quad f^3_{46} = f^4_{63} = -\frac{1}{2q}.$$
(2.19)

If $p \neq q$ one can choose the following coordinate representation³ for the one-forms (e^i, ω^a) :

$$\begin{aligned} e^1 &= -\sin \psi d\theta_1 + \sin \theta_1 \cos \psi d\phi_1, \\ e^2 &= \cos \psi d\theta_1 + \sin \theta_1 \sin \psi d\phi_1, \\ e^3 &= -\sin \psi d\theta_2 - \sin \theta_2 \cos \psi d\phi_2, \\ e^4 &= -\cos \psi d\theta_2 + \sin \theta_2 \sin \psi d\phi_2, \\ e^5 &= -[(p-q)d\psi + p \cos \theta_1 d\phi_1 + q \cos \theta_2 d\phi_2], \\ \omega^1 &= -[(p+q)d\psi + p \cos \theta_1 d\phi_1 - q \cos \theta_2 d\phi_2], \end{aligned}$$
(2.20)

where $(\theta_{1,2}, \phi_{1,2})$ are spherical coordinates on $\mathbb{CP}^1_{A,B}$ respectively, satisfying $0 \leq \theta_{1,2} < \pi$, $0 \leq \phi_{1,2} < 2\pi$, and ψ describes the $U(1)$ -bundle, with $0 \leq \psi < 2\pi$.

From eq. (2.10) describing the action of the $U(1)$ we find that the case of interest is $p = -q = 1$. One finds then from (2.16) that in the coframe basis e^i , a left-invariant metric should satisfy

$$g_{11} = g_{22} = m_1, \quad g_{33} = g_{44} = m_2, \quad g_{55} = m_3, \quad g_{13} = g_{24} = m_4, \quad g_{14} = -g_{23} = m_5,$$
(2.21)

with $(m_1, m_2, m_3, m_4, m_5)$ five functions of the transversal coordinate. In the explicit representation (2.20) the metric becomes

$$\begin{aligned} ds^2 &= m_1 [(d\theta_1)^2 + \sin^2 \theta_1 (d\phi_1)^2] + m_2 [(d\theta_2)^2 + \sin^2 \theta_2 (d\phi_2)^2] + \\ &\quad m_3 [2d\psi + \cos \theta_1 d\phi_1 - \cos \theta_2 d\phi_2]^2 \\ &\quad + (-m_4 \cos 2\psi + m_5 \sin 2\psi)(d\theta_1 d\theta_2 + \sin \theta_1 \sin \theta_2 d\phi_1 d\phi_2) \\ &\quad + (m_4 \sin 2\psi + m_5 \cos 2\psi)(\sin \theta_2 d\theta_1 d\phi_2 - \sin \theta_1 d\theta_2 d\phi_1). \end{aligned}$$
(2.22)

³For $p = q$ this particular representation is not so great, since the terms with $d\psi$ drop out of e^5 . For this case another more suitable coordinate representation can be found by changing some signs. Here we are interested in the case $p = -q = 1$ anyway.

Furthermore, from (2.15) one finds that the left-invariant forms are spanned by

$$\begin{aligned}
&1 - \text{forms} : e^5, \\
&2 - \text{forms} : e^{12}, e^{34}, e^{14} - e^{23}, e^{13} + e^{24}, \\
&3 - \text{forms} : e^{125}, e^{345}, e^{145} - e^{235}, e^{135} + e^{245}, \\
&4 - \text{forms} : e^{1234}, \\
&5 - \text{forms} : e^{12345}.
\end{aligned} \tag{2.23}$$

Finally, we remark that the structure constants (2.19) are invariant under a shift of ψ :

$$\psi \rightarrow \psi + \phi(t), \tag{2.24}$$

where we can allow the shift to depend explicitly on the transversal coordinate t . This implies that the conditions we will impose on our solutions in section 3 are invariant under this shift, which will thus send $\text{SO}(4)$ -invariant solutions to $\text{SO}(4)$ -invariant solutions. If ϕ is constant, the shift reduces to the $\text{SO}(2)_R$ symmetry (2.12). The non-constant part we will appropriately gauge away.

2.3 Starting point: the massless Kähler-Einstein geometry

The starting point of the analysis is the $N = 6$ $\text{SU}(4)$ -invariant vacuum on $\mathbb{CP}^3$ with the standard Fubini-Study metric in terms of the homogeneous coordinates (Z^1, Z^2, Z^3, Z^4) . I will rewrite this vacuum in terms of the coordinates induced by the local foliation with $T^{1,-1}$ leaves. Only the $\text{SO}(4)$ isometry group and $N = 2$ supersymmetry are then manifest.

Corresponding to the description of $T^{1,-1}$ as a $\text{U}(1)$ -bundle over $\mathbb{CP}^1 \times \mathbb{CP}^1$, appropriate coordinates are

$$z = \frac{Z^1}{Z^2}, \quad w = \frac{Z^3}{Z^4}, \quad \psi = \frac{1}{2} \arg \left(\frac{Z^1 Z^2}{Z^3 Z^4} \right), \tag{2.25}$$

where z and w are inhomogeneous coordinates for $\mathbb{CP}_{A,B}^1$ respectively. Furthermore it will be convenient to choose a transversal coordinate ξ as follows

$$\tan^2 \xi = t = \frac{|Z^1|^2 + |Z^2|^2}{|Z^3|^2 + |Z^4|^2}, \tag{2.26}$$

and thus $0 \leq \xi \leq \frac{\pi}{2}$. If we further replace the inhomogeneous coordinates (z, w) for both $\mathbb{CP}^1$ s by the respective spherical coordinates $(\theta_{1,2}, \phi_{1,2})$ we find in the end:

$$\begin{aligned}
Z^1 &= \lambda \sin \xi \cos \frac{\theta_1}{2} \exp \frac{i}{2} (\psi + \phi_1), \\
Z^2 &= \lambda \sin \xi \sin \frac{\theta_1}{2} \exp \frac{i}{2} (\psi - \phi_1), \\
Z^3 &= \lambda \cos \xi \cos \frac{\theta_2}{2} \exp \frac{i}{2} (-\psi + \phi_2), \\
Z^4 &= \lambda \cos \xi \sin \frac{\theta_2}{2} \exp \frac{i}{2} (-\psi - \phi_2),
\end{aligned} \tag{2.27}$$

where λ is the inessential overall complex factor of the homogeneous coordinates.

The Fubini-Study metric becomes in these coordinates:

$$a^{-1}ds^2 = d\xi^2 + \frac{\sin^2 \xi}{4} [(d\theta_1)^2 + \sin^2 \theta_1 (d\phi_1)^2] + \frac{\cos^2 \xi}{4} [(d\theta_2)^2 + \sin^2 \theta_2 (d\phi_2)^2] \\ + \sin^2 \xi \cos^2 \xi \left(d\psi + \frac{1}{2} \cos \theta_1 d\phi_1 - \frac{1}{2} \cos \theta_2 d\phi_2 \right)^2, \quad (2.28)$$

where we introduced a constant overall scale $a > 0$.

Note that the orbits indeed correspond to $T^{p,q}$ with $p = -q = 1$, and furthermore to putting

$$m_1 = \frac{\sin^2 \xi}{4}, \quad m_2 = \frac{\cos^2 \xi}{4}, \quad m_3 = \frac{\sin^2 \xi \cos^2 \xi}{4}, \quad m_4 = m_5 = 0, \quad (2.29)$$

in the coset metric of (2.22). The transversal coordinate ξ is chosen so that $g_{\xi\xi} = a$ is constant. These coordinates for $\mathbb{CP}^3$ were already proposed in [39].

Since it is just the standard Fubini-Study metric in new coordinates, we know that this metric is regular on the degenerate orbits $\xi \rightarrow 0$ and $\xi \rightarrow \pi/2$, where respectively the first and the second $\mathbb{CP}^1$ as well as the fiber collapse. It is a useful exercise to check this explicitly, since it shows us what to pay attention to when imposing the boundary conditions on the general solutions. In the limit $\xi \rightarrow 0$ the metric becomes

$$a^{-1}ds^2 = d\xi^2 + \frac{\xi^2}{4} [(d\theta_1)^2 + \sin^2 \theta_1 (d\phi_1)^2 + (2d\psi + \cos \theta_1 d\phi_1 - \cos \theta_2 d\phi_2)^2] \\ + \frac{1}{4} [(d\theta_2)^2 + \sin^2 \theta_2 (d\phi_2)^2]. \quad (2.30)$$

For constant (ξ, θ_2, ϕ_2) the second term in the first line becomes the standard metric of an S^3 . Checking the volume of this $S^3(\xi)$ we find:

$$\text{vol}(S^3(\xi)) = \left(\frac{\xi}{2}\right)^3 \int \sin \theta_1 d\theta_1 d\phi_1 2d\psi = (2\pi^2)\xi^2, \quad (2.31)$$

the standard volume of an S^3 with radius ξ . For this calculation the prefactor $1/4$ in front of the second term in the first line, together with the period of ψ is crucial. It follows that the part of the metric in the first line is just the flat $\mathbb{R}^4$ metric, and thus regular. In a completely analogous way, the metric is also regular for $\xi \rightarrow \pi/2$.

The standard closed Kähler form J' is in these coordinates given by

$$a^{-1}J' = \frac{\sin^2 \xi}{4} e^{12} + \frac{\cos^2 \xi}{4} e^{34} + \frac{\sin 2\xi}{4} e^5 \wedge d\xi. \quad (2.32)$$

Note that this is *not* the two-form J of the $SU(3)$ -structure associated to either of the supersymmetries. In particular, J' is only associated to a $U(3)$ -structure – the one in the denominator of the coset description (2.3) – since it will induce an integrable complex structure of which the $(3,0)$ -form is globally only well-defined up to a factor. The existence of these two two-forms, J' and J , and the associated integrable and non-integrable complex structures is a generic property of twistor spaces [42, 43]. See also [1] for a discussion of this in the context of the homogeneous $N = 1$ type IIA solutions. We will construct the two-forms associated to the $SU(3)$ -structures in the next section after we review the conditions these $SU(3)$ -structures have to satisfy.

3. Supersymmetry conditions and Bianchi identities

3.1 Review of the conditions for supersymmetric AdS₄ solutions

In this subsection we review the conditions for supersymmetric strict SU(3)-structure AdS₄ flux compactifications, first derived in [8]. These conditions are equivalent to the set of supersymmetry conditions and Bianchi identities without sources, which as we mentioned in the introduction suffices to imply all the remaining equations of motion.

It is found that the internal manifold should possess an SU(3)-structure. This consists of a real two-form J and a complex decomposable three-form Ω satisfying the compatibility and normalization condition

$$\Omega \wedge J = 0, \quad (3.1a)$$

$$\Omega \wedge \bar{\Omega} = \frac{4i}{3} J^3 \neq 0, \quad (3.1b)$$

and such that the associated metric is positive-definite. Especially the condition for Ω to be complex decomposable is quite complicated and was studied in [44]. In fact, it was shown in that paper that this implies that $\text{Im}\Omega$ is (up to a sign) determined by $\text{Re}\Omega$. This works as follows: from $\text{Re}\Omega$ we can construct first an almost complex structure. We define

$$\tilde{\mathcal{I}}_k{}^l = \varepsilon^{lm_1\dots m_5} (\text{Re}\Omega)_{km_1m_2} (\text{Re}\Omega)_{m_3m_4m_5}, \quad (3.2)$$

where $\varepsilon^{m_1\dots m_6}$ is the epsilon-tensor in six dimensions, and then properly normalize it

$$\mathcal{I} = \frac{\tilde{\mathcal{I}}}{\sqrt{-\text{tr} \frac{1}{6} \tilde{\mathcal{I}}^2}}, \quad (3.3)$$

so that $\mathcal{I}^2 = -\mathbb{1}$.

$$H(\text{Re}\Omega) = \text{tr} \frac{1}{6} \tilde{\mathcal{I}}^2 \quad (3.4)$$

is called the Hitchin function. This procedure only works if the Hitchin function is strictly negative, which imposes a condition on $\text{Re}\Omega$. The metric can then be constructed from $\mathcal{I}$ and J through

$$g_{mn} = \mathcal{I}_m{}^l J_{ln}, \quad (3.5)$$

and $\text{Im}\Omega$ is given by

$$\text{Im}\Omega = \mp \frac{1}{3} \mathcal{I}_k{}^l dx^k \wedge \iota_l \text{Re}\Omega. \quad (3.6)$$

Furthermore, the only non-zero SU(3)-structure torsion classes are the scalar $\mathcal{W}_1$ and the primitive (1,1)-form $\mathcal{W}_2$, so that

$$dJ = -\frac{3}{2} i \mathcal{W}_1 \text{Re}\Omega, \quad (3.7a)$$

$$d\Omega = \mathcal{W}_1 J \wedge J + \mathcal{W}_2 \wedge J. \quad (3.7b)$$

They can be chosen to be both purely imaginary, and the sourceless Bianchi identity for F_2 imposes moreover

$$d\mathcal{W}_2 = ic_2 \text{Re}\Omega, \quad (3.8)$$

where from the defining properties of $\mathcal{W}_2$ we find for the proportionality factor

$$c_2 = -\frac{1}{8}|\mathcal{W}_2^-|^2. \quad (3.9)$$

The warp factor A and the dilaton Φ are constant, and the Romans mass is given by

$$\frac{16}{5}e^{2\Phi}m^2 = 3|\mathcal{W}_1|^2 - |\mathcal{W}_2|^2 \geq 0, \quad (3.10)$$

imposing an inequality on the torsion classes. Finally the RR- and NSNS fluxes are then given by

$$H = \frac{2m}{5}e^\Phi \text{Re}\Omega, \quad (3.11a)$$

$$F_2 = \frac{f}{9}J + F'_2, \quad (3.11b)$$

$$F_4 = f\text{vol}_4 + \frac{3m}{10}J \wedge J, \quad (3.11c)$$

where we have defined

$$e^\Phi f = \frac{9i}{4}\mathcal{W}_1, \quad e^\Phi F'_2 = i\mathcal{W}_2. \quad (3.12)$$

Massaging these conditions a bit and keep only the essential, in the end we have to solve the following problem. Find a geometry with two-forms $J, \widetilde{\mathcal{W}}_2, L$ and a three-form $\text{Re}\Omega$ so that

$$dJ = c_1 \text{Re}\Omega, \quad (3.13a)$$

$$\widetilde{\mathcal{W}}_2 = \frac{\tilde{c}_2}{c_1}J + L, \quad dL = 0, \quad (3.13b)$$

$$J \wedge \text{Re}\Omega = 0, \quad L \wedge \text{Re}\Omega = 0, \quad (3.13c)$$

$$\text{Re}\Omega \wedge \text{Im}\Omega(\text{Re}\Omega) = -\frac{2}{3}J^3 \neq 0, \quad (3.13d)$$

$$d[\text{Im}\Omega(\text{Re}\Omega)] = \widetilde{\mathcal{W}}_2 \wedge J, \quad (3.13e)$$

where the notation $\text{Im}\Omega(\text{Re}\Omega)$ stresses that $\text{Im}\Omega$ should be found from $\text{Re}\Omega$ through the procedure outlined in eqs. (3.2-3.6), making the conditions containing $\text{Im}\Omega$ the most involved. After we find a solution to the above conditions we must still check for positivity of the metric. The relation with the parameters introduced earlier is as follows

$$\begin{aligned} c_1 &= -\frac{2}{3}e^\Phi f, \\ \widetilde{\mathcal{W}}_2 &= \frac{2}{3}c_1 J - i\mathcal{W}_2, \\ \tilde{c}_2 &= \frac{2}{3}(c_1)^2 + c_2, \\ e^{2\Phi}m^2 &= \frac{5}{4}(2\tilde{c}_2 - (c_1)^2) = \frac{5}{4}(2\rho - 1)(c_1)^2 \geq 0. \end{aligned} \quad (3.14)$$

We introduced the parameter

$$\rho = \frac{\tilde{c}_2}{(c_1)^2}, \quad (3.15)$$

which considering $m^2 \geq 0$ and $c_2 \leq 0$ obeys $\frac{1}{2} \leq \rho \leq \frac{2}{3}$. The limiting cases correspond to the massless and nearly-Kähler case respectively.

Since we want to construct a solution that respects the $\text{SO}(4)$ isometry, we expand the unknown forms in left-invariant forms, which we build from the forms of (2.23) and the extra left-invariant one-form $d\xi$,

$$\begin{aligned} J &= k_1(\xi)e^{12} + k_2(\xi)e^{34} + k_3(\xi)d\xi \wedge e^5 + k_4(\xi) [\cos \theta_k(\xi)(e^{13} + e^{24}) + \sin \theta_k(\xi)(e^{14} - e^{23})], \\ \text{Re}\Omega &= u_1(\xi)e^{125} + u_2(\xi)e^{345} + u_3(\xi)(e^{145} - e^{235}) + u_4(\xi)(e^{135} + e^{245}) + \\ &\quad u_5(\xi)d\xi \wedge e^{12} + u_6(\xi)d\xi \wedge e^{34} + u_7(\xi)d\xi \wedge (e^{14} - e^{23}) + u_8(\xi)d\xi \wedge (e^{13} + e^{24}) \\ L &= l_1(\xi)e^{12} + l_2(\xi)e^{34} + l_3(\xi)d\xi \wedge e^5 + l_4(\xi)(e^{14} - e^{23}) + l_5(\xi)(e^{13} + e^{24}), \end{aligned} \quad (3.16)$$

where the introduction of the angle coordinate $\theta_k(\xi)$ will be convenient later on, and we do not display $\widehat{\mathcal{W}}_2$ since it can be trivially expressed in terms of J and L through (3.13b).

Before solving these conditions in general, we will first use them to find the $\text{SU}(3)$ -structures corresponding to the two $\text{SO}(4)$ -invariant supersymmetries of the massless Kähler-Einstein geometry.

3.2 The massless Kähler-Einstein geometry revisited

In this case we already know the metric explicitly and that such J and $\text{Re}\Omega$, corresponding to the two supersymmetries invariant under $\text{SO}(4)$, exist. We just want to find them in the coordinate system of section 2.3. It turns out that to find most of J (i.e. apart from $\theta_k(\xi)$) we do not have to solve the full set of conditions (3.13). It suffices to plug the ansatz for J , (3.16), into (3.5), and further impose $\mathcal{I}^2 = -\mathbb{1}$ (for a proper complex structure) and $J \wedge dJ = 0$ (which follows from (3.13a) and (3.13c)). The other conditions then become relatively simple, and in the end we find for J and Ω :

$$\begin{aligned} a^{-1}J &= -\frac{\sin^2 \xi \cos 2\xi}{4}e^{12} + \frac{\cos^2 \xi \cos 2\xi}{4}e^{34} + \frac{\sin 2\xi}{4}d\xi \wedge e^5 \\ &\quad + \frac{\sin^2 2\xi}{8} [\cos \theta_k (e^{13} + e^{24}) + \sin \theta_k (e^{14} - e^{23})] \\ a^{-3/2}\Omega &= \left(d\xi - \frac{i}{4} \sin 2\xi e^5 \right) \wedge \frac{1}{2} [\cos^2 \xi (ie^3 + e^4) - \sin^2 \xi e^{-i\theta_k} (e^1 - ie^2)] \\ &\quad \wedge \frac{1}{4} \sin 2\xi [ie^{i\theta_k} (e^1 + ie^2) + e^3 + ie^4], \end{aligned} \quad (3.17)$$

where $\theta_k(\xi) = \theta_k$ is constant and we chose the upper sign in (3.6). We find furthermore

$$c_1 = \frac{4}{\sqrt{a}}, \quad \tilde{c}_2 = \frac{1}{2}(c_1)^2. \quad (3.18)$$

Note that in general we can get three more solutions by using one or both of

$$\begin{aligned} J &\rightarrow J, \quad \text{Re}\Omega \rightarrow -\text{Re}\Omega, \quad \text{Im}\Omega \rightarrow -\text{Im}\Omega, \quad c_1 \rightarrow -c_1, \quad \tilde{c}_2 \rightarrow \tilde{c}_2, \\ J &\rightarrow -J, \quad \text{Re}\Omega \rightarrow \text{Re}\Omega, \quad \text{Im}\Omega \rightarrow -\text{Im}\Omega, \quad c_1 \rightarrow -c_1, \quad \tilde{c}_2 \rightarrow \tilde{c}_2, \end{aligned} \quad (3.19)$$

where the second line corresponds to choosing a different sign in (3.6). In the following we will always choose $c_1 > 0$ and the upper sign in (3.6).

The angle θ_k corresponds to the freedom of rotating the $SU(3)$ -structure with $SO(2)_R$. In more detail, applying the transformation (2.12) on the solution results in $\theta_k \rightarrow \theta_k + 2\phi$. It does not change the metric, nor, since $m = 0$ any of the fluxes⁴. It follows that the solution has two $SO(4)$ -invariant supersymmetries.

Furthermore we remark that the coset $T^{1,-1}$ is invariant under

$$(e^1, e^2) \leftrightarrow (e^3, e^4), \quad e^5 \rightarrow -e^5, \quad (3.20)$$

which is completed with

$$\xi \rightarrow \frac{\pi}{2} - \xi, \quad \theta_k \rightarrow -\theta_k, \quad (3.21)$$

into a symmetry of the solution, which, as we will find, extends to the general solution.

Finally, it will be convenient later on to use the freedom of reparameterizing ξ to put $k_4(\tilde{\xi}) = \tilde{\xi}$. For the $m = 0$ solution above we find then

$$\tilde{\xi} = \frac{a}{8} \sin^2 2\xi, \quad k_1(\tilde{\xi})k_2(\tilde{\xi}) = \tilde{\xi}^2 - \frac{a}{8}\tilde{\xi}, \quad k_1(\tilde{\xi}) + k_2(\tilde{\xi}) = \frac{a}{4} - 2\tilde{\xi}. \quad (3.22)$$

3.3 Solving the conditions

In this section, we put ourselves to the task of solving the equations (3.13) for general $1/2 \leq \rho \leq 2/3$. As it turns out, the equations (3.13a)-(3.13c) are relatively easily to solve and put:

$$\begin{aligned} u_1(\xi) &= u_2(\xi) = l_4(\xi) = l_5(\xi) = 0, \\ k_3(\xi) &= \frac{[k_1(\xi)k_2(\xi) - k_4(\xi)^2]'}{k_1(\xi) - k_2(\xi)}, \\ l_3(\xi) &= -l_1'(\xi), \quad l_2(\xi) = d_1 - l_1(\xi), \\ l_1(\xi) &= d_1 \frac{k_1(\xi)[k_1(\xi) + k_2(\xi)]' - [k_4(\xi)^2]'}{[k_1(\xi) + k_2(\xi)][k_1(\xi) + k_2(\xi)]' - 2[k_4(\xi)^2]'}, \\ u_3(\xi) &= -\frac{k_4(\xi) \cos \theta_k(\xi)}{c_1}, \quad u_4(\xi) = \frac{k_4(\xi) \sin \theta_k(\xi)}{c_1}, \\ u_5(\xi) &= \frac{k_1(\xi)[k_1(\xi) + k_2(\xi)]' - [k_4(\xi)^2]'}{c_1[k_1(\xi) - k_2(\xi)]}, \quad u_6(\xi) = -\frac{k_2(\xi)[k_1(\xi) + k_2(\xi)]' - [k_4(\xi)^2]'}{c_1[k_1(\xi) - k_2(\xi)]}, \\ u_7(\xi) &= \frac{[k_4(\xi) \sin \theta_k(\xi)]'}{c_1}, \quad u_8(\xi) = \frac{[k_4(\xi) \cos \theta_k(\xi)]'}{c_1}, \end{aligned} \quad (3.23)$$

where prime denotes the derivative to ξ . This leaves four unknown functions $k_1(\xi)$, $k_2(\xi)$, $k_4(\xi)$, $\theta_k(\xi)$, and one unknown constant d_1 . It turns out that none of the equations in (3.13) puts a constraint on $\theta_k(\xi)$. This follows from the fact that (2.24) is a symmetry of the equations and the solutions with different $\theta_k(\xi)$ are related to each other by this shift.

⁴This is somewhat less obvious for F_2 , but can still be checked using (3.11,3.12).

We will gauge-fix the non-constant part of this coordinate reparametrization by requiring $g_{\xi\psi}$ to vanish, which forces $\theta_k(\xi) = \theta_k$ to be constant.

To proceed it will be convenient to also fix the reparametrization freedom of ξ by introducing a new coordinate $\tilde{\xi}$ such that $k_4(\tilde{\xi}) = \tilde{\xi}$. Furthermore, if we introduce

$$p(\tilde{\xi}) = \tilde{\xi}^2 - k_1(\tilde{\xi})k_2(\tilde{\xi}), \quad s(\tilde{\xi}) = k_1(\tilde{\xi}) + k_2(\tilde{\xi}), \quad (3.24)$$

and square the left- and right-hand side of (3.13d) – removing the square root in (3.3) – it takes the following simplified form

$$\frac{\tilde{\xi}^2 \left\{ [s(\tilde{\xi}) - \tilde{\xi}s'(\tilde{\xi})]^2 + p(\tilde{\xi})[4 - s'(\tilde{\xi})^2] \right\}}{\left\{ \frac{1}{2} [p(\tilde{\xi})^2] \right\}'^2} = (c_1)^4. \quad (3.25)$$

This is still not easy to solve unless $s'(\tilde{\xi})^2 = 4$, which is indeed the case for the massless solution (3.22). We will first make this ansatz, and discuss the alternative case afterwards.

The ansatz $s'(\tilde{\xi})^2 = 4$

More specifically, we put

$$s(\tilde{\xi}) = e_1 - 2\tilde{\xi}, \quad (3.26)$$

where we made the same choice of sign as in (3.22). There is no loss of generality in this sign choice, since it can be changed using (3.19). We can then solve (3.25) and find

$$p(\tilde{\xi})^2 = \frac{e_1}{(c_1)^2} \tilde{\xi}^2 + e_2. \quad (3.27)$$

Finally we turn to condition (3.13e). It should hold when we wedge it with the most general left-invariant two-form, which contains five degrees of freedom. Because we have already imposed that $\widetilde{\mathcal{W}}_2$ is a (1,1)-form, it is automatically satisfied when wedging with a (2,0)- or (0,2)-form. This removes two degrees of freedom. One more degree of freedom is eliminated because (3.13e) is independent of $\theta_k(\xi)$. So we need only impose

$$d\text{Im}\Omega \wedge L = \widetilde{\mathcal{W}}_2 \wedge L \wedge J \iff \widetilde{\mathcal{W}}_2 \wedge L \wedge J = 0, \quad (3.28a)$$

$$d\text{Im}\Omega \wedge J = \widetilde{\mathcal{W}}_2 \wedge J \wedge J \iff \widetilde{\mathcal{W}}_2 \wedge J \wedge J = \frac{2}{3} c_1 J^3. \quad (3.28b)$$

Of course, we found it prudent to check the full (3.13e) after finding the solutions.

Plugging in (3.26) and (3.27) we find in the end two sets of solutions in terms of ρ

$$e_2 = 0, \quad e_1 = \frac{20 \mp 8\sqrt{4-6\rho} - 6\rho}{(c_1)^2(2+\rho)^2}, \quad d_1 = \frac{-4 \pm 4\sqrt{4-6\rho} + 6\rho}{c_1(2+\rho)}. \quad (3.29)$$

Let us now transform back to the original coordinates where $g_{\xi\xi}$ is constant. We need then to solve for

$$\sqrt{g_{\xi\xi}(\tilde{\xi})} d\tilde{\xi} = a^{1/2} d\xi. \quad (3.30)$$

Further normalizing such that $0 \leq \xi \leq \pi/2$ we find

$$\tilde{\xi} = \frac{c_1(e_1)^{3/2}}{4(c_1\sqrt{e_1}-1)} \sin^2 2\xi, \quad a = \frac{4e_1}{c_1\sqrt{e_1}-1}. \quad (3.31)$$

The resulting metric is given by

$$\begin{aligned} a^{-1}ds^2 = & d\xi^2 + \frac{\sin^2 \xi}{8} [c_1\sqrt{e_1} + (2 - c_1\sqrt{e_1}) \cos 2\xi] [(d\theta_1)^2 + \sin^2 \theta_1 (d\phi_1)^2] \\ & + \frac{\cos^2 \xi}{8} [c_1\sqrt{e_1} - (2 - c_1\sqrt{e_1}) \cos 2\xi] [(d\theta_2)^2 + \sin^2 \theta_2 (d\phi_2)^2] \\ & + \sin^2 \xi \cos^2 \xi \left(d\psi + \frac{1}{2} \cos \theta_1 d\phi_1 - \frac{1}{2} \cos \theta_2 d\phi_2 \right)^2 \\ & + \frac{\sin^2 2\xi}{16} (2 - c_1\sqrt{e_1}) \left[\sin(2\psi + \theta_k) (d\theta_1 d\theta_2 + \sin \theta_1 \sin \theta_2 d\phi_1 d\phi_2) \right. \\ & \left. + \cos(2\psi + \theta_k) (\sin \theta_2 d\theta_1 d\phi_2 - \sin \theta_1 d\theta_2 d\phi_1) \right]. \end{aligned} \quad (3.32)$$

It is easy to check that for $\xi = \epsilon$ and $\xi = \pi/2 - \epsilon$ with ϵ infinitesimal, the first three lines of the metric take exactly the same form as the metric in the massless Kähler-Einstein case in these limits, while the terms in the last two lines vanish as ϵ^2 . The metric is thus regular.

Except for $2 - c_1\sqrt{e_1} = 0$, which is the case for the massless Kähler-Einstein solution, the metric changes when shifting θ_k . Moreover the fluxes will also change. So the $SO(2)_R$ symmetry is broken, although it still sends solutions to solutions.

Furthermore, we find that the $SU(3)$ -structure is given by

$$\begin{aligned} J(\theta_k) = & \frac{1}{4(c_1\sqrt{e_1}-1)} \left\{ 2e_1 \sin^2 \xi (-2 + c_1\sqrt{e_1} - c_1\sqrt{e_1} \cos 2\xi) e^{12} \right. \\ & + 2e_1 \cos^2 \xi (-2 + c_1\sqrt{e_1} + c_1\sqrt{e_1} \cos 2\xi) e^{34} + 4e_1 \sin 2\xi d\xi \wedge e^5 \\ & \left. + c_1(e_1)^{3/2} \sin^2 2\xi [\cos \theta_k (e^{13} + e^{24}) + \sin \theta_k (e^{14} - e^{23})] \right\}, \end{aligned} \quad (3.33)$$

$$\Omega(\theta_k) = \frac{(c_1)^3(e_1)^{3/2}}{8(c_1\sqrt{e_1}-1)} \Omega_{KE}(\theta_k),$$

where $\Omega_{KE}(\theta_k)$ is the holomorphic three-form of the Kähler-Einstein case defined in (3.17). The fact that Ω is proportional to Ω_{KE} means that the complex structure is the same on all solutions with the same θ_k .

Beyond the ansatz

We will now show that there are no further solutions beyond the ansatz $s'(\tilde{\xi})^2 = 4$. An important role is played by the regularity of the metric for $\tilde{\xi} \rightarrow 0$. In terms of the unknown functions $p(\tilde{\xi})$ and $s(\tilde{\xi})$ the relevant components of the metric are:

$$\begin{aligned} g_{\tilde{\xi}\tilde{\xi}} = & \frac{(c_1)^2 p(\tilde{\xi}) p'(\tilde{\xi})^2}{\tilde{\xi}^2 (-4\tilde{\xi}^2 + s(\tilde{\xi})^2 + 4p(\tilde{\xi}))}, \quad g_{55} = \frac{\tilde{\xi}^2}{(c_1)^2 p(\tilde{\xi})}, \\ g_{11}, g_{33} = & \frac{\tilde{\xi}}{2(c_1)^2 p(\tilde{\xi}) p'(\tilde{\xi})} \left[\left(s(\tilde{\xi}) - \tilde{\xi} s'(\tilde{\xi}) \right) \left(s(\tilde{\xi}) \mp \sqrt{-4\tilde{\xi}^2 + s(\tilde{\xi})^2 + 4p(\tilde{\xi})} \right) + 4p(\tilde{\xi}) \right], \end{aligned} \quad (3.34)$$

where the upper/lower sign is for g_{11}, g_{33} respectively. Requiring that — after an appropriate coordinate transformation $\xi(\tilde{\xi})$ such that $g_{\xi\xi} = 1$ — the metric takes the form (2.30) for $\xi \rightarrow 0$, we find for the asymptotic behavior of $p(\tilde{\xi})$ and $s(\tilde{\xi})$:

$$p(\tilde{\xi}) = d_1 \tilde{\xi} + \mathcal{O}(\tilde{\xi}^2), \quad s(\tilde{\xi}) = d_2 + \mathcal{O}(\tilde{\xi}), \quad (3.35)$$

with $(d_2)^2 = (c_1)^4 (d_1)^4$. Solving now eqs. (3.25) and (3.28) order by order in $\tilde{\xi}$ we recover solution (3.26), (3.27), (3.29).

3.4 Equivalence to the family of $N = 1$ massive $\text{Sp}(2)$ -invariant solutions

In this subsection we show that the family of solutions (3.32), (3.33) is equivalent to the family of $N = 1$ $\text{Sp}(2)$ -invariant solutions of [1, 17]. It will be convenient to use the procedure discussed in section 5.1 of [29] to construct the $\text{Sp}(2)$ -invariant solutions in terms of the homogeneous coordinates on $\mathbb{CP}^3$.

As in that paper we start with $\mathbb{C}^4 \setminus \{0\}$, which can be considered as a $\mathbb{C}^*$ bundle over $\mathbb{CP}^3$ with projection map p . We can describe the coordinates Z^a of $\mathbb{C}^4$ in terms of our $\text{SO}(4)$ -adapted coordinates as in (2.27), where we take $\lambda = r e^{i\gamma}$. The fiber coordinates are then (r, γ) . We are interested in forms α on the total space of the bundle $\mathbb{C}^4 \setminus \{0\}$ that are the pull-back under p of a form on the base space $\mathbb{CP}^3$. Such forms are called *basic*. A basic form is both *vertical*, i.e. $\iota_v \alpha = 0$ for all vectors v along the fiber, and *invariant*, i.e. $\mathcal{L}_v \alpha = 0$. For our bundle, the vectors tangent to the fiber are spanned by

$$r \partial_r = Z^a \partial_a + \bar{Z}_a \bar{\partial}^a, \quad \partial_\gamma = i(Z^a \partial_a - \bar{Z}_a \bar{\partial}^a), \quad (3.36)$$

and the dual one-forms are

$$\frac{dr}{r} = \frac{1}{2r^2} (\bar{Z}_a dZ^a + Z^a d\bar{Z}_a), \quad \eta = \frac{i}{2r^2} (-\bar{Z}_a dZ^a + Z^a d\bar{Z}_a). \quad (3.37)$$

One can now split a form on $\mathbb{C}^4$ into a vertical and a non-vertical part. For the one-forms dZ^a of the complex basis, for instance, one finds:

$$dZ^a = \left(\delta^a_b - \frac{Z^a \bar{Z}_b}{r^2} \right) dZ^b + Z^a \left(\frac{dr}{r} + i\eta \right). \quad (3.38)$$

Using this one finds immediately for the standard Kähler form on $\mathbb{C}^4$,

$$J_{(4)} = \frac{i}{2} dZ^a \wedge d\bar{Z}_a = r dr \wedge \eta + r^2 J_{\text{FS}}, \quad J_{\text{FS}} = \frac{i}{2r^2} DZ^a D\bar{Z}_a, \quad (3.39)$$

where J_{FS} is the unit Fubini-Study Kähler form. One can indeed check that J_{FS} is vertical and invariant. Changing to the $\text{SO}(4)$ -adapted coordinates (2.27) it takes the form $a^{-1} J'$ of (2.32). Using $\mathcal{L}_{r\partial_r} J_{(4)} = 2J_{(4)}$, which follows from the fact that $J_{(4)}$ is homogeneous of degree two in the coordinates Z^a , we find furthermore:

$$d\eta = J_{\text{FS}}. \quad (3.40)$$

On the other hand, if one decomposes

$$\Omega_{(4)} = dZ^1 \wedge dZ^2 \wedge dZ^3 \wedge dZ^4 = -ir^3 (dr + ir\eta) \wedge \Omega_{\text{FS}}, \quad (3.41)$$

one finds that Ω_{FS} is vertical and satisfies $\mathcal{L}_{r\partial_r}\Omega_{\text{FS}} = 0$. However, Ω_{FS} is not basic since $\mathcal{L}_\gamma\Omega_{\text{FS}} = 4i\Omega_{\text{FS}} \neq 0$. This is related to the fact that on $\mathbb{CP}^3$ there is no globally defined $(3,0)$ -form associated to J_{FS} .

Let us now make a choice of $\text{Sp}(2)$ subgroup of $\text{SU}(4)$. Such a choice is in one-to-one correspondence with the choice of a *holomorphic symplectic form* κ . This is a closed two-form, $d\kappa = 0$, whose square gives the holomorphic $(4,0)$ -form

$$\frac{1}{2}\kappa^2 = \Omega_{(4)} . \quad (3.42)$$

Indeed, just like the supersymmetry generators, κ transforms as a **6** under $\text{SU}(4)$, which decomposes under $\text{Sp}(2)$ as $\mathbf{6} \rightarrow \mathbf{1} + \mathbf{5}$. κ then corresponds to the invariant part. Expanding κ in a vertical and a non-vertical part, one obtains a two-form t_κ and a one-form s_κ as follows

$$\kappa = r(dr + ir\eta) \wedge s_\kappa + r^2 t_\kappa . \quad (3.43)$$

By construction, these forms are vertical and one also easily checks their invariance under $r\partial_r$. They are however *not* invariant under ∂_γ . The simplest way to obtain this, is to first realize that $\mathcal{L}_{r\partial_r}\kappa = 2\kappa$, since κ is homogeneous of degree two in the complex coordinates Z^a . Together with (3.43) one finds then

$$ds_\kappa = 2(i\eta \wedge s_\kappa + t_\kappa) , \quad (3.44)$$

and finally, using (3.40),

$$\mathcal{L}_\gamma s_\kappa = 2is_\kappa , \quad \mathcal{L}_\gamma t_\kappa = 2it_\kappa , \quad (3.45)$$

so that s_κ and t_κ are *not* well-defined on $\mathbb{CP}^3$. From (3.42) it follows that, in fact, Ω_{FS} , which is also not well-defined on $\mathbb{CP}^3$, can be written as

$$\Omega_{\text{FS}} = is_\kappa \wedge t_\kappa . \quad (3.46)$$

However, it is now obvious that if we define instead

$$\Omega = i\bar{s}_\kappa \wedge t_\kappa , \quad (3.47)$$

the γ -charges compensate and we do obtain a well-defined $(3,0)$ -form on $\mathbb{CP}^3$. This corresponds to a different choice of almost complex structure, replacing one one-form of the complex basis by its conjugate. If we want (J, Ω) to still correspond to the Fubini-Study metric we have to define J by performing the same operation, i.e. $s_\kappa \leftrightarrow \bar{s}_\kappa$, on J_{FS} . This amounts to⁵

$$J = -(J_{\text{FS}} - is_\kappa \wedge \bar{s}_\kappa) . \quad (3.48)$$

In fact, generalizing this construction we can define a whole family of different $\text{SU}(3)$ -structures $(J_\sigma, \Omega_\sigma)$, corresponding to different metrics on $\mathbb{CP}^3$, as follows:

$$\begin{aligned} J_\sigma &= -a \left(\frac{2}{\sigma} J_{\text{FS}} - i \frac{\sigma + 2}{2\sigma} s_\kappa \wedge \bar{s}_\kappa \right) , \\ \Omega_\sigma &= \frac{2ia^{3/2}}{\sigma} \bar{s}_\kappa \wedge t_\kappa , \end{aligned} \quad (3.49)$$

⁵The overall sign in the definition of J ensures that $J^3 = J_{\text{FS}}^3$.

where $a > 0$ is an overall scale and σ is a shape parameter. One can check that these $(J_\sigma, \Omega_\sigma)$ solve the supersymmetry conditions (3.7) and (3.8) with

$$-i\mathcal{W}_1 = \frac{2c_1}{3} = \frac{2(2+\sigma)}{3a^{1/2}}, \quad |\mathcal{W}_2|^2 = -8c_2 = \frac{64(1-\sigma)^2}{3a}. \quad (3.50)$$

The condition (3.10) is satisfied for $2/5 \leq \sigma \leq 2$, and the fluxes can be found from (3.11) and (3.12). The parameter ρ of eq. (3.15) is given by

$$\rho = \frac{2\sigma(4-\sigma)}{(2+\sigma)^2}. \quad (3.51)$$

Since these solutions also manifestly preserve the $\text{Sp}(2)$ singled out by κ , they correspond in fact to the $N = 1$ family of solutions first introduced in [1].

From the discussion of section 2.1 it follows that to make contact with our solutions we should take

$$\kappa = e^{i\theta/2} dZ^1 \wedge dZ^2 + e^{-i\theta/2} dZ^3 \wedge dZ^4. \quad (3.52)$$

Indeed, our specific choice of $\text{SO}(4)$, of which the action was defined around (2.5), leaves both terms in (3.52) separately invariant, corresponding to $\mathbf{6} \rightarrow (\mathbf{1}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{1}) \oplus (\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2})$. So any choice of θ corresponds to a choice of $\text{Sp}(2)$ so that $\text{SO}(4) \subset \text{Sp}(2)$. Going through the above construction, we find that $(J_\sigma, \Omega_\sigma)$ exactly matches (3.33) upon putting

$$e_1 = \frac{a}{4}, \quad \theta_k = \theta - \frac{\pi}{2}. \quad (3.53)$$

The transformation (2.12) corresponds to shifting $\theta \rightarrow \theta + 2\phi$. It will rotate the $\text{SU}(3)$ -structure $(J_\sigma, \Omega_\sigma)$ and thus supersymmetry generator as well as the $\text{Sp}(2)$ symmetry group. In the massless Kähler-Einstein case, $\sigma = 2$, it leaves the metric and the fluxes invariant so that it really corresponds to an R-symmetry, while in all other cases it rotates a solution into a different solution.

We conclude that the solutions of (3.33) enhance their symmetry group to $\text{Sp}(2)$ and form the $N = 1$ family of homogeneous solutions first introduced in [1]. Moreover, we have shown that there are no other solutions with $\text{SO}(4)$ symmetry group and strict $\text{SU}(3)$ -structure.

4. Coisotropic D-brane embeddings

Now that we have constructed the $\text{SU}(3)$ -structures associated to the supersymmetry in an $\text{SO}(4)$ -invariant description, let us see whether the geometry allows for the embedding of supersymmetric D-branes, which according to [31, 32] must be generalized calibrated. It is known [26, 27, 28] that there are special Lagrangian D-branes wrapping $\mathbb{RP}^3$ s, but we will here be interested in the more exotic coisotropic D8-branes [34].

A coisotropic space-time-filling D8-brane has one transversal direction in $\mathbb{CP}^3$ and is thus a priori compatible with the symmetry group $\text{SO}(4)$, acting with cohomogeneity one. So let us construct an $\text{SO}(4)$ -invariant coisotropic D8-brane. To be left-invariant, the one-form part of the D-brane source j , defining the direction transversal to the D-brane, must

be a linear combination of the one-forms $d\xi$ and e^5 . In fact, the only coordinate dependence allowed by the $SO(4)$ symmetry group is the dependence on ξ , so if we want a localized D-brane, we must take the transversal coordinate to be ξ . The coisotropic D-brane then wraps $T^{1,-1}$ at some constant value $\xi = \xi_0$. For the world-volume gauge field $\mathcal{F}$ we make the $SO(4)$ -invariant ansatz

$$\mathcal{F} = f_1 e^{12} + f_2 e^{34} + f_3 (e^{14} - e^{23}) + f_4 (e^{13} + e^{24}) , \quad (4.1)$$

with f_i constant. It must satisfy

$$d\mathcal{F} = H|_{\Sigma} . \quad (4.2)$$

The generalized calibration condition [31, 32] for a space-time-filling D8-brane becomes

$$(iJ|_{\Sigma} + \mathcal{F})^2 = 0 , \quad \text{Re}\Omega|_{\Sigma} \wedge \mathcal{F} = 0 . \quad (4.3)$$

The first condition, which can be interpreted as an F-flatness condition in the low-energy effective theory [45], is equivalent to the condition for a coisotropic D-brane [34], see e.g. [35, 46]. This has to be completed with the second condition, which can be interpreted as a D-flatness condition. However, in [33] it was shown that in AdS_4 compactifications the second condition follows automatically from the first.

Let us first consider the massless Kähler-Einstein background. In this case, we find from $H = 0$ that $\mathcal{F}$ must be closed, implying $f_3 = f_4 = 0$. Furthermore, we find from (4.3)

$$\xi_0 = \pi/4, \quad 64 f_1 f_2 = -a^2, \quad \text{or} \quad \xi_0 = 0, \quad f_1 = 0, \quad \text{or} \quad \xi_0 = \pi/2, \quad f_2 = 0, \quad (4.4)$$

Since (4.3) then holds no matter the choice of θ in (3.17), the D-brane preserves both $SO(4)$ -invariant supersymmetries. Only the first possibility leads to a genuine D8-brane, since for the last two possibilities the second respectively the first $\mathbb{CP}^1$ and the fiber shrink to zero and the D-brane only wraps the other $\mathbb{CP}^1$.

In the full quantum theory, the closed world-volume gauge field $F_{\text{WV}} = \mathcal{F} - B|_{\Sigma}$ must be quantized

$$\frac{1}{(l_s)^2} \int_{S^2} F_{\text{WV}} = n_F , \quad (4.5)$$

where $l_s = 2\pi\sqrt{\alpha'}$ is the string length, S^2 is a representative of the two-cycle homology of $T^{1,-1}$ and n_F is an integer. We can take $B|_{\Sigma} = 0$. This will boil down to

$$\frac{a}{(l_s)^2} d(\tilde{f}_1 + \tilde{f}_2) = n_F , \quad (4.6)$$

where d is a coefficient of order one (depending on the details of the integral) and we introduced the dimensionless $\tilde{f}_i = a^{-1} f_i$. There is no problem in choosing the $\tilde{f}_i$ such that n_F is integer and $64 \tilde{f}_1 \tilde{f}_2 = -1$ (which follows from (4.4)).

We now move on to the general massive $N = 1$ background (3.33), where we put (3.51) and (3.53). This time, since $H \neq 0$ the condition on the exterior derivative of $\mathcal{F}$ is more complicated:

$$d\mathcal{F} = H|_{\Sigma} = \frac{2m}{5} e^{\Phi} \text{Re}\Omega|_{\Sigma} = \sqrt{\frac{-5\sigma^2 + 12\sigma - 4}{5a}} \text{Re}\Omega|_{\Sigma} , \quad (4.7)$$

where we used (3.11a) and (3.10). The calibration conditions are still the same as in (4.3). We find the following solution:

$$\begin{aligned}
f_1 &= a \sin^2 \xi_0 \frac{\sigma(2+\sigma)\sqrt{-5\sigma^2+12\sigma-4} \sin^2 2\xi_0 \mp 8\sqrt{\sigma^3[1-3\sigma-(1+2\sigma)\cos 4\xi_0]}}{8\sqrt{5}\sigma^2[-2+\sigma+(2+\sigma)\cos 2\xi_0]}, \\
f_2 &= a \cos^2 \xi_0 \frac{\sigma(2+\sigma)\sqrt{-5\sigma^2+12\sigma-4} \sin^2 2\xi_0 \pm 8\sqrt{\sigma^3[1-3\sigma-(1+2\sigma)\cos 4\xi_0]}}{8\sqrt{5}\sigma^2[-2+\sigma-(2+\sigma)\cos 2\xi_0]}, \\
f_3 &= a \frac{\sqrt{-5\sigma^2+12\sigma-4}}{16\sqrt{5}\sigma} \sin^2 2\xi_0 \sin \theta_k, \\
f_4 &= a \frac{\sqrt{-5\sigma^2+12\sigma-4}}{16\sqrt{5}\sigma} \sin^2 2\xi_0 \cos \theta_k,
\end{aligned} \tag{4.8}$$

where the D-brane can be at any $\xi = \xi_0$ such that $1 - 3\sigma - (1 + 2\sigma)\cos 4\xi_0 \geq 0$ or $\frac{1}{4} \arccos\left(\frac{1-3\sigma}{1+2\sigma}\right) \leq \xi_0 \leq \frac{\pi}{2} - \frac{1}{4} \arccos\left(\frac{1-3\sigma}{1+2\sigma}\right)$. The D-brane then preserves the $N = 1$ supersymmetry of the background. In the Kähler-Einstein limit, $\sigma \rightarrow 2$, the allowed interval shrinks to just the point $\xi_0 = \frac{\pi}{4}$ and we obtain the $N = 2$ solution above. Moving towards the squashed massless solution, $\sigma \rightarrow 2/5$, on the other hand, the interval gets larger and becomes $\frac{1}{4} \arccos\left(-\frac{1}{9}\right) \leq \xi_0 \leq \frac{\pi}{2} - \frac{1}{4} \arccos\left(-\frac{1}{9}\right)$ at this point.

Using the freedom in ξ_0 , again solutions can be found such that the world-volume gauge field is quantized as in (4.5). This time the B -field is non-zero, but it can still be globally defined. We can take $B = \frac{2m}{5c_1} \text{Re}\Omega$. This B -field also plays an important role in the flux quantization of the bulk RR-charges (for a discussion see [1, 21]). Indeed, the RR-fields have to be twisted with e^B in order to obtain closed forms with well-defined periods.

5. Concluding remarks

In this paper I have shown that the only solutions on $\text{AdS}_4 \times \mathbb{CP}^3$ with symmetry group $\text{SO}(4)$ and a strict $\text{SU}(3)$ -structure ansatz for the supersymmetry, are the homogeneous $N = 1$ solutions of [1], which possess the larger symmetry group $\text{Sp}(2)$. This is in agreement with the results of [2], where although a Chern-Simons-matter theory with $\text{SO}(4)$ global symmetry was proposed, it was also argued that the geometric dual should have $\text{SU}(3) \times \text{SU}(3)$ -structure instead. In the meantime, in [29] this dual has been constructed to first order in the Romans mass parameter m , making use of the explicit form of the D2-brane superpotential, obtained from the CFT. This is particularly interesting since, so far, there were no examples of source-less supergravity compactifications with dynamic $\text{SU}(3) \times \text{SU}(3)$ -structure (see [30] for another recent construction). The construction of the full solution seems more difficult, since the $\text{SU}(3) \times \text{SU}(3)$ supersymmetry conditions are more complicated than the ones for the strict $\text{SU}(3)$ -structure ansatz. The basic procedure should however be the same as in this paper, namely to expand the pure spinors defining the $\text{SU}(3) \times \text{SU}(3)$ -structure in $\text{SO}(4)$ left-invariant forms and try to solve the differential equations in one variable. As in [29] one could also insert the extra data of the D2-brane superpotential.

Furthermore, I have presented probe supersymmetric D8-branes of the coisotropic type on both the Kähler-Einstein ABJM geometry as well as the massive $\mathrm{Sp}(2)$ family. It would be interesting to find the dual CFT that would correspond to adding D8-branes in such a way. Presumably, as is the case for adding special Lagrangian D6-branes [26, 27, 28] this would add flavors to the Chern-Simons-matter theory.

Finally, one could also try to construct similar solutions on the $\frac{\mathrm{SU}(3)}{\mathrm{U}(1) \times \mathrm{U}(1)}$ manifold, for which e.g. [47] could provide a starting point.

Acknowledgments

I would like to thank Claudio Caviezel, Johanna Erdmenger, Simon Körs and Luca Martucci for useful discussions. Furthermore I thank Dario Martelli and James Sparks for pointing out a fatal flaw in the first version, and Davide Gaiotto and Alessandro Tomasiello for suggesting an approach to rewrite the paper. For many of the calculations I found it convenient to use Mathematica together with the package `scalarEDCcode` for handling forms (<http://library.wolfram.com/infocenter/MathSource/683>).

I am supported by the German Research Foundation (DFG) within the Emmy-Noether-Program (Grant number ZA 279/1-2).

References

- [1] A. Tomasiello, *New string vacua from twistor spaces*, *Phys. Rev.* **D78** (2008) 046007, [[arXiv:0712.1396](#)].
- [2] D. Gaiotto and A. Tomasiello, *The gauge dual of Romans mass*, [arXiv:0901.0969](#).
- [3] M. Graña, *Flux compactifications in string theory: A comprehensive review*, *Phys. Rept.* **423** (2006) 91–158, [[hep-th/0509003](#)].
- [4] M. R. Douglas and S. Kachru, *Flux compactification*, *Rev. Mod. Phys.* **79** (2007) 733–796, [[hep-th/0610102](#)].
- [5] R. Blumenhagen, B. Körs, D. Lüst, and S. Stieberger, *Four-dimensional string compactifications with D-branes, orientifolds and fluxes*, *Phys. Rept.* **445** (2007) 1–193, [[hep-th/0610327](#)].
- [6] F. Denef, *Les Houches lectures on constructing string vacua*, [arXiv:0803.1194](#).
- [7] J. P. Gauntlett and S. Pakis, *The geometry of $D = 11$ Killing spinors*, *JHEP* **04** (2003) 039, [[hep-th/0212008](#)].
- [8] D. Lüst and D. Tsimpis, *Supersymmetric AdS_4 compactifications of IIA supergravity*, *JHEP* **02** (2005) 027, [[hep-th/0412250](#)].
- [9] J. P. Gauntlett, D. Martelli, J. Sparks, and D. Waldram, *Supersymmetric AdS_5 solutions of type IIB supergravity*, *Class. Quant. Grav.* **23** (2006) 4693–4718, [[hep-th/0510125](#)].
- [10] P. Koerber and D. Tsimpis, *Supersymmetric sources, integrability and generalized-structure compactifications*, *JHEP* **08** (2007) 082, [[arXiv:0706.1244](#)].
- [11] J. M. Maldacena and C. Núñez, *Supergravity description of field theories on curved manifolds and a no go theorem*, *Int. J. Mod. Phys.* **A16** (2001) 822–855, [[hep-th/0007018](#)].

- [12] O. Aharony, O. Bergman, D. L. Jafferis, and J. Maldacena, $\mathcal{N} = 6$ superconformal Chern-Simons-matter theories, M2-branes and their gravity duals, *JHEP* **10** (2008) 091, [[arXiv:0806.1218](#)].
- [13] K. Behrndt and M. Cvetič, General $\mathcal{N} = 1$ supersymmetric flux vacua of (massive) type IIA string theory, *Phys. Rev. Lett.* **95** (2005) 021601, [[hep-th/0403049](#)].
- [14] K. Behrndt and M. Cvetič, General $\mathcal{N} = 1$ supersymmetric fluxes in massive type IIA string theory, *Nucl. Phys.* **B708** (2005) 45–71, [[hep-th/0407263](#)].
- [15] T. House and E. Palti, Effective action of (massive) IIA on manifolds with $SU(3)$ structure, *Phys. Rev.* **D72** (2005) 026004, [[hep-th/0505177](#)].
- [16] G. Aldazabal and A. Font, A second look at $\mathcal{N} = 1$ supersymmetric AdS_4 vacua of type IIA supergravity, *JHEP* **02** (2008) 086, [[arXiv:0712.1021](#)].
- [17] P. Koerber, D. Lüst, and D. Tsimpis, Type IIA AdS_4 compactifications on cosets, interpolations and domain walls, *JHEP* **07** (2008) 017, [[arXiv:0804.0614](#)].
- [18] B. E. W. Nilsson and C. N. Pope, Hopf fibration of eleven-dimensional supergravity, *Class. Quant. Grav.* **1** (1984) 499.
- [19] C. Caviezel, P. Koerber, S. Körs, D. Tsimpis, D. Lüst, and M. Zagermann, The effective theory of type IIA AdS_4 compactifications on nilmanifolds and cosets, *Class. Quant. Grav.* **26** (2009) 025014, [[arXiv:0806.3458](#)].
- [20] S. Körs, On the effective theory of type IIA AdS_4 compactifications, [arXiv:0810.5104](#).
- [21] D. Cassani and A.-K. Kashani-Poor, Exploiting $\mathcal{N}=2$ in consistent coset reductions of type IIA, *Nucl. Phys.* **B817** (2009) 25–57, [[arXiv:0901.4251](#)].
- [22] H. Ooguri and C.-S. Park, Superconformal Chern-Simons theories and the squashed seven sphere, *JHEP* **11** (2008) 082, [[arXiv:0808.0500](#)].
- [23] M. Fujita, W. Li, S. Ryu, and T. Takayanagi, Fractional quantum Hall effect via holography: Chern-Simons, edge states, and hierarchy, [arXiv:0901.0924](#).
- [24] D. Lüst, F. Marchesano, L. Martucci, and D. Tsimpis, Generalized non-supersymmetric flux vacua, *JHEP* **11** (2008) 021, [[arXiv:0807.4540](#)].
- [25] D. Lüst and D. Tsimpis, Classes of AdS_4 type IIA/IIB compactifications with $SU(3) \times SU(3)$ structure, *JHEP* **04** (2009) 111, [[arXiv:0901.4474](#)].
- [26] S. Hohenegger and I. Kirsch, A note on the holography of Chern-Simons matter theories with flavour, [arXiv:0903.1730](#).
- [27] D. Gaiotto and D. L. Jafferis, Notes on adding D6 branes wrapping $\mathbb{RP}^3$ in $AdS_4 \times \mathbb{CP}^3$, [arXiv:0903.2175](#).
- [28] Y. Hikida, W. Li, and T. Takayanagi, ABJM with Flavors and FQHE, [arXiv:0903.2194](#).
- [29] D. Gaiotto and A. Tomasiello, Perturbing gauge/gravity duals by a Romans mass, [arXiv:0904.3959](#).
- [30] M. Petrini and A. Zaffaroni, $\mathcal{N} = 2$ solutions of massive type IIA and their Chern-Simons duals, [arXiv:0904.4915](#).
- [31] P. Koerber, Stable D-branes, calibrations and generalized Calabi-Yau geometry, *JHEP* **08** (2005) 099, [[hep-th/0506154](#)].

- [32] L. Martucci and P. Smyth, *Supersymmetric D-branes and calibrations on general $N = 1$ backgrounds*, *JHEP* **11** (2005) 048, [[hep-th/0507099](#)].
- [33] P. Koerber and L. Martucci, *D-branes on AdS flux compactifications*, *JHEP* **01** (2008) 047, [[arXiv:0710.5530](#)].
- [34] A. Kapustin and D. Orlov, *Remarks on A-branes, mirror symmetry, and the Fukaya category*, *J. Geom. Phys.* **48** (2003) 84, [[hep-th/0109098](#)].
- [35] A. Font, L. E. Ibáñez, and F. Marchesano, *Coisotropic D8-branes and model-building*, *JHEP* **09** (2006) 080, [[hep-th/0607219](#)].
- [36] A. Sevrin, W. Staessens, and A. Wijns, *An $N = 2$ worldsheet approach to D-branes in bihermitian geometries: I. Chiral and twisted chiral fields*, *JHEP* **10** (2008) 108, [[arXiv:0809.3659](#)].
- [37] A. Kapustin and E. Witten, *Electric-magnetic duality and the geometric Langlands program*, [hep-th/0604151](#).
- [38] P. Candelas and X. C. de la Ossa, *Comments on Conifolds*, *Nucl. Phys.* **B342** (1990) 246–268.
- [39] M. Cvetič, H. Lü, and C. N. Pope, *Consistent warped-space Kaluza-Klein reductions, half-maximal gauged supergravities and $\mathbb{CP}_n$ constructions*, *Nucl. Phys.* **B597** (2001) 172–196, [[hep-th/0007109](#)].
- [40] F. Müller-Hoissen and R. Stückl, *Coset spaces and ten-dimensional unified theories*, *Class. Quant. Grav.* **5** (1988) 27.
- [41] P. van Nieuwenhuizen, *General theory of coset manifolds and antisymmetric tensors applied to Kaluza-Klein supergravity*, in *Supersymmetry and supergravity '84*, World Scientific, Singapore, 1985.
- [42] M. F. Atiyah, H. N.J., and S. I.M., *Self-duality in four-dimensional riemannian geometry*, *Proc. Roy. Soc. London Ser. A* **362** (1978) 425–461.
- [43] J. Eels and S. Salamon, *Constructions twistorielles des applications harmoniques*, *C.r. Acad. sc. Paris, sér. I Math.* **296** (1983) 685–687.
- [44] N. Hitchin, *The geometry of three-forms in six and seven dimension*, [math/0010054](#).
- [45] L. Martucci, *D-branes on general $N = 1$ backgrounds: Superpotentials and D-terms*, *JHEP* **06** (2006) 033, [[hep-th/0602129](#)].
- [46] P. Koerber and L. Martucci, *Deformations of calibrated D-branes in flux generalized complex manifolds*, *JHEP* **12** (2006) 062, [[hep-th/0610044](#)].
- [47] D. L. Jafferis and A. Tomasiello, *A simple class of $\mathcal{N} = 3$ gauge/gravity duals*, *JHEP* **10** (2008) 101, [[arXiv:0808.0864](#)].